



# Meat products and functional food

Slaviša Stajić<sup>a\*</sup>, Nikola Stanišić<sup>b</sup> and Vladimir Kurćubić<sup>c</sup>

<sup>a</sup> University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Belgrade, Serbia

<sup>b</sup> Institute for Animal Husbandry, Autoput 16, 11080 Belgrade, Serbia

<sup>c</sup> University of Kragujevac, Faculty of Agronomy, Cara Dušana 34, 32000 Čačak, Serbia

## ARTICLE INFO

### Keywords:

Meat products  
Functional food  
Sodium reduction  
Additive replacement  
n-3 fatty acids  
Dietary fibres

## ABSTRACT

Functional food is a term for food products fortified with ingredients that possess beneficial physiological effects. Meat products are characterized by relatively high salt, fat and saturated fatty acid contents. Moreover, phosphates and nitrites are also marked as potential harmful components. Designing meat products as functional food has been associated with replacement (and/or reduction) of these components with other, especially natural, ingredients that possess beneficial effects. The development of such products poses quite a challenge since it requires the creation of a product with improved functional properties and the same sensory quality as conventional ones. Results of numerous studies into improving the nutritional properties of meat products indicate that the meat industry has responded to the changes of lifestyle and perception of food.

## 1. Introduction

Meat is generally a significant source of several nutrients: proteins of high biological value, and micronutrients such as iron, zinc, phosphorus, selenium and vitamin B12 (Pereira & Vicente, 2013; Williams, 2007). Fat is a great source of energy. Moreover, meat is almost an exclusive source of several bioactive compounds with antioxidative, anti-inflammatory, anti-carcinogenic and anti-atherosclerotic properties, such as conjugated linoleic acids, carnosine, anserine and taurine (Pereira & Vicente, 2013; Young *et al.*, 2013). Since meat as essential food needed to be preserved, ancient people developed meat products by combining different animal tissues with salt and spices, and applying early preservation techniques — drying, heating and smoking (Kurćubić *et al.*, 2022). For centuries, these products have been an excellent source of protein, energy and other nutri-

ents. However, in the last 50–60 years, fresh meat has become readily available and meat products lost their primary function and are more valuable because of their sensory properties (Stajić & Vasilev, 2022).

In the last third of the 20<sup>th</sup> century, it was observed that some food ingredients can have a negative effect on health (salt, saturated fatty acids (SFA)) if they are consumed in sufficient quantities. Conversely, other ingredients (antioxidants, n-3 fatty acids, minerals, vitamins) can be important in preventing or treating certain diseases (Doyon & Labrecque, 2008). This led to a different perception of food — the purpose of food is no longer only to satisfy hunger and provide energy and basic nutrients, but it could also be a tool that prevents the occurrence of diseases caused by changes in lifestyle and diet, and improves physical and mental health (Siró *et al.*, 2008). Therefore, the concept of “new food” appeared which later became “functional food”.

\*Corresponding author: Stajić Slaviša, [stajic@agrif.bg.ac.rs](mailto:stajic@agrif.bg.ac.rs)

Paper received Jun 9<sup>th</sup> 2023. Paper accepted Jun 18<sup>th</sup> 2023.

Published by Institute of Meat Hygiene and Technology — Belgrade, Serbia

This is an open access article under CC BY licence (<http://creativecommons.org/licenses/by/4.0>)

## 2. Functional food

In 1984, Japanese scientists first used the term “functional food” for food products fortified with ingredients that possess beneficial physiological effects (Bigliardi & Galati, 2013). Japan is also the first country where this field was regulated (in 1991), when the Ministry of Health introduced the regulation called FOSHU — Food for Specified Health Uses (Iwatani & Yamamoto, 2019). Since then, numerous definitions of functional food have been proposed — Bigliardi and Galati (2013) selected 39 definitions after extensive literature research. In general, functional food should be food that is part of regular nutrition and contains ingredients with a selective effect on one or more functions of the body, and the positive effects of which can be seen as physiologically functional (Jiménez-Colmenero *et al.* 2010; Zhang *et al.*, 2010).

## 3. Meat products as functional food

In general, meat products contain all the nutrients/compounds from meat that have a positive effect on health, although the content of some minerals, taurine, carnosine and thermolabile vitamins, can be reduced during the process of production, storage and preparation for consumption. Some compounds with a positive effect on health (e.g. bioactive peptides) and others with an adverse effect on health (e.g. biogenic amines and nitrosamines) can occur during the production process. Also, some ingredients such as salt (sodium), phosphates and nitrites, as well as SFA that fatty tissue is rich with, are correlated with a negative influence on health. A large number of research studies has investigated reduction and/or replacement of these potential harmful components.

### 3.1. Reduction of salt (sodium)

Salt is the most common non-meat ingredient used in meat processing. It is essential for technological and sensory properties, shelf-life and safety of meat products. Its influence depends on salt content, form of meat (whole muscle/ground), processing procedures (heating, drying, mixing/grinding) and can be summarized as: extraction and activation of myofibrillar proteins, meat emulsion formulation and stabilization, improvement of the water-holding capacity (WHC) and defining flavour (Kurćubić *et al.*, 2022). High sodium intake has been corre-

lated with high blood pressure, while salt is the major source of sodium in meat products, in which it accounts for approximately 79% of the sodium (Desmond, 2006).

The salt content in s formulation depends on the type of meat product: usually 1–2% in burger type products, 1.8–2.5% in emulsion-type sausages, 2–3% in cooked ham, 2.5–3% in dry-fermented sausages (>3.5% after drying), >3% in dry-cured meats (>4.5% after drying). The literature data indicate that salt reduction to about 1.7% in emulsion-type sausages (without phosphates added) and to 2.3% in fermented sausages can be reached without major adverse effects on products (Corral *et al.*, 2013; Stajić *et al.*, 2022). The major impact of salt on WHC came from the Cl<sup>-</sup> anion (Petit *et al.*, 2019); therefore, reduction of the salt content in meat products has limitations. For example, salt levels above 1.3% (with added phosphates) are needed to obtain a stable meat emulsion (Vasquez Mejia *et al.*, 2019). Another possible course of action is the reduction of the sodium content by replacing salt with other chloride salts (KCl, CaCl<sub>2</sub>, MgCl<sub>2</sub>) which also has limitations, because the salty taste of salt mainly comes from the Na<sup>+</sup> cation (Petit *et al.*, 2019), and the use of other chloride salts can alter taste — e.g. use of KCl and MgCl<sub>2</sub> leads to bitterness (Corral *et al.*, 2013). Therefore, the partial replacement of NaCl with other chloride salts and combining it with compounds/ingredients that enhance salty taste and/or mask bitter taste could be good solution (Inguglia *et al.*, 2017). Different commercial mixtures consisting of sodium salt mixtures and lysine, arginine, K-lactate, glycine, yeast and mushroom extracts have been developed with the aim of reducing the sodium level and overcoming/masking technological and sensory defects (Kurćubić *et al.*, 2022).

### 3.2. Replacement of phosphates

Phosphates are very important for meat products because they increase the swelling of meat fibres (WHC), promote solubilization of myofibrillar proteins, bind metal ions, and reduce viscosity of meat batters (Sebranek, 2009). Therefore, phosphates are of great significance for technological properties of emulsified-type sausages and whole-muscle brine-injected meat products. The reduction of the phosphate content was not in focus (as much as salt and fat reduction) because the content/addition of phosphates to meat products is limited by regulations — 8 g/kg product of total phosphorus (as

P<sub>2</sub>O<sub>5</sub>) in Serbia, and 5 g/kg product of added phosphates in the EU. However, due to a change in P/Ca ratio in dietary intake and the increase of consumer demand for the “clean label” products, several research studies have been conducted to examine phosphate replacement, especially in emulsion-type sausages. The phosphate reduction was based on: i) replacement with natural mineral-based ingredients e.g. calcium powders originating from shells, eggs and algae (Bae et al., 2017; Stajić et al., 2020); ii) replacement with natural ingredients that can bind water and emulsify fat, e.g. dietary fibre and heteropolysaccharides (Câmara et al., 2020; Powell et al., 2019; Stajić et al., 2022); iii) reduction by application of alternative processing techniques e.g. ultrasound (Pinton et al., 2019).

Natural calcium powders increase pH values and, therefore, contribute to proper processing yield, while on the other hand, they alter colour and texture properties (Bae et al., 2017; Stajić et al., 2020). Dietary fibre contains soluble and insoluble fibre and so has gel-forming ability, water-binding capacity and oil-binding capacity (Tunland & Meyer, 2002). However, due to differences in the amount of soluble and insoluble fibre, different impacts occur that are amount- and product-dependent. Lower emulsion stability was obtained when phosphates were replaced with 0.3–0.6% of wheat, maize, pea, potato and 0.5–1% of citrus fibre (Powell et al., 2019; Stajić et al., 2022). However, this was not significant in all treatments, and similar processing yields compared to controls were obtained in treatments with 0.6% of wheat, maize and pea fibres and 0.75% of citrus fibre. Application of ultrasound for 18 minutes can be used in processing emulsion-type sausages with 50% of reduced phosphates. These strategies have certain limitations; however, they offer great potential to multi-ingredient phosphate replacement strategy.

### 3.3. Nitrate replacement/reduction

Nitrites are ingredients that have multiple effects in meat systems: antimicrobial effects (particularly on neurotoxin-producing *Clostridium botulinum*), pink colour and aroma formation and lipid oxidation delay (Alirezalu et al., 2019). However, nitrites also participate in the creation of carcinogenic N-nitrosamines, whose content is in correlation with the content of nitrites (Alirezalu et al., 2019). Nitrites are usually added in amounts of 100–150 mg/kg depending on regulations. The main part of this amount is needed for *C. botulinum* con-

trol, while about 25 mg is needed for colour formation (Sindelar & Milkowski, 2012). Therefore, the strategy for nitrite reduction (and thus N-nitrosamine reduction) includes the introduction of ingredients that exhibit antimicrobial and antioxidant activity. In the research of Kurćubić et al. (2014), nitrite was replaced with ethanol extract of *Kitabeilia vitifolia* in effective concentration of 12.5 g/kg of the initial batch of dry-fermented sausages. The results indicate the great antimicrobial and antioxidative potential of *K. vitifolia* ethanol extract during production and cold storage. Also, colour, taste and overall acceptability were not affected. In frankfurter-type sausages, Alirezalu et al. (2019) obtained promising results in terms of antimicrobial and antioxidant effects and sensory properties of frankfurter-type sausages during cold storage (45 days at 4°C) when replacing nitrite with 1% of chitosan and 0.2% of ε-polylysine (both in combination with a 500 ppm mixture of green tea, stinging nettle and olive leaves extracts).

### 3.4. Fat reduction and/or improvement of fatty acids profile

Fatty tissue is essential for the quality of meat products. This is especially significant in meat products where fatty tissue is ground together with meat, mixed with non-meat ingredients (salt, additives, spices), and subjected to different procedures (drying, fermentation, grinding, emulsification) to produce fermented sausages, emulsion-type sausages or burgers/patties (Kurćubić et al., 2022). The fat content of these products can be up to 50% in dry-fermented sausages and up to 30% in emulsion-type sausages and burgers/patties (Kurćubić et al., 2022). Moreover, animal fat is rich in SFA and has a low content of n-3 polyunsaturated fatty acids (PUFA). Recommendations of the World Health Organization (WHO) from two decades ago emphasized that the amount of fat in total daily energy intake should be in the 15–30% range, SFA < 10% and n-3 PUFA 1–2%. Strategies include reduction of the amount of fatty tissue and/or partial to total replacement with non-lipid replacers or oils rich in PUFA (Kurćubić et al., 2022). The reduction of the amount of fatty tissue has limited effects because this reduces the acceptability of fermented sausages (Liaros et al., 2009) and burgers (Heck et al., 2019). A partial replacement of fatty tissue with non-lipid fat replacers (inulin, cereal, and fruit fibre) can be a strategy to improve the quality of low-fat meat products

(Bajcic *et al.*, 2023; Kurćubić *et al.*, 2020). However, this only reduces the intake of energy which originates from fat — the FA profile can be changed only by introducing oils rich in PUFA (especially n-3 PUFA) into the formulation of meat products (Kurćubić *et al.*, 2022). On the other hand, oils rich in PUFA (e.g. grapeseed, flaxseed, fish, algae and their combinations) are more susceptible to lipid oxidation, and therefore, these oils need to be stabilized (cannot be added in liquid form) before application (Stajić & Vasilev, 2022). Oils were immobilized and stabilized in emulsions, double emulsions, gel-like matrixes (hydrogel, oleogel, oil-bulking, structured emulsions) and encapsulated by different encapsulation techniques — spray-drying, electrostatic extrusion, etc. (Stajić & Vasilev, 2022). The immobilization technique together with oil type, amount of fat replacement, and the principle of fat replacement (with same amount of oil (later stabilized) or with the same amount of substitute which consists of immobilized oil) influence the technological and sensory properties of modified meat products (Stajić *et al.*, 2018; Stajić *et al.*, 2020). Nutritional properties depend on the amount of fat replacement and oil fatty acid profile.

### 3.5. Meat products as sources of dietary fibre

As mentioned above, food could also be a tool that prevents the occurrence of diseases. Meat products are not a source of dietary fibre, however, so could hybrid meat products, made by combining two or more types of meat with plant proteins and other non-meat ingredients, have found their place on the market (Galanakis *et al.*, 2021). Hybrid meat products have a potential advantage over plant-based meats, because they provide the familiar meaty taste and texture while containing plant-based ingredients that can contribute to a healthier and sustainable diet (Grasso & Javorska, 2020). They can offer a broader range of flavours and textures while reportedly having signif-

icantly lower greenhouse gas emissions when compared to all-meat products (Baune *et al.*, 2021).

The percentage of plant ingredients (legumes, grains, fruits and vegetables) in these products can be from 10% to 50%. By definition, they are not added as extensions but as part of the product constituents (Grasso & Javorska, 2020). Ingredients like soy, wheat, starch and fibre have been used in the meat industry for their functional properties (emulsification, water binding/holding and gelling) and to save costs (Singh *et al.*, 2008; Asgar *et al.*, 2010). The conceptual difference between hybrid and traditional meat products is that plant-based ingredients are used not only for economic and technological purposes, but also for improving health claims, lowering the environmental impact and decreasing meat consumption (Grasso, 2020; Talens *et al.*, 2022). A market research study showed that a third of consumers are flexitarians, which means that although they eat meat, they regularly avoid it on certain days (Grasso and Javorska, 2020). Therefore, hybrid meat products could create new business opportunities for the meat industry, as they fulfil the growing flexitarian consumer needs.

## 4. Conclusion

Results of numerous studies into improving the nutritional properties of meat products indicate that the meat industry has responded to the changes of lifestyle and perceptions of the food. Meat products with reduced sodium and fat contents and improved fatty acid profiles have been developed. Moreover, natural replacers for synthetic and potentially harmful ingredients have been introduced. The development of such products poses quite a challenge since it requires the creation of a product with improved functional properties and the same sensory quality as conventional ones. New research is being carried out with the aim of optimizing the developed products in order to provide the necessary sensory quality in addition to further improving the nutritional properties.

**Disclosure statement:** No potential conflict of interest was reported by the authors.

**Funding:** This work was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia, project ref. number: 451-03-47/2023-01/200116.

## References

- Alirezalu, K., Hesari, J., Nemati, Z., Munekata, P. E. S., Barba, F. J. & Lorenzo, J. M. (2019). Combined effect of natural antioxidants and antimicrobial compounds during refrigerated storage of nitrite-free frankfurter-type sausage. *Food Research International*, 120, 839–850, <https://doi.org/10.1016/j.foodres.2018.11.048>
- Asgar, M. A., Fazilah, A., Huda, N., Bhat, R. & Karim, A. A. (2010). Nonmeat protein alternatives as meat extenders and meat analogs. *Comprehensive Reviews in Food Science & Food Safety*, 9, 513–529, <http://dx.doi.org/10.1111/j.1541-4337.2010.00124.x>
- Bae, S. M., Cho, M. G. & Jeong, J. Y. (2017). Effects of various calcium powders as replacers for synthetic phosphate on the quality properties of ground pork meat products. *Korean Journal for Food Science of Animal Resources*, 37, 456–463, <http://dx.doi.org/10.5851/kosfa.2017.37.3.456>
- Bajcic, A., Petronijevic, R., Suvajdzic, B., Tomovic, V., Stajković, S. & Vasilev, D. (2023). Use of inulin-collagen suspension for the total replacement of pork backfat in cooked-emulsified sausages. *Journal of Food and Nutrition Research*, 62, 35–45.
- Barros, J. C., Munekata, P. E. S., de Carvalho, F. A. L., Domínguez, R., Trindade, M. A., Pateiro, M. & Lorenzo, J. M. (2021). Healthy beef burgers: Effect of animal fat replacement by algal and wheat germ oil emulsions. *Meat Science*, 173, 108396, <https://doi.org/10.1016/j.meatsci.2020.108396>
- Baune, M.-C., Jeske, A.-L., Profeta, A., Smetana, S., Broucke, K., Van Royen, G., Gibis, M., Weiss, J. & Terjung, N. (2021). Effect of plant protein extrudates on hybrid meatballs — changes in nutritional composition and sustainability. *Future Foods*, 4, 100081, <https://doi.org/10.1016/j.fufo.2021.100081>
- Bigliardi, B. & Galati, F. (2013). Innovation trends in the food industry: The case of functional foods. *Trends in Food Science & Technology*, 31, 118–129, <http://dx.doi.org/10.1016/j.tifs.2013.03.006>
- Câmara, A. K. F. I., Vidal, V. A. S., Santos, M., Bernardinelli, O. D., Sabadini, E. & Pollonio, M. A. R. (2020). Reducing phosphate in emulsified meat products by adding chia (*Salvia hispanica* L.) mucilage in powder or gel format: A clean label technological strategy. *Meat Science*, 163, 108085, <https://doi.org/10.1016/j.meatsci.2020.108085>
- Corral, S., Salvador, A. & Flores, M. (2013). Salt reduction in slow fermented sausages affects the generation of aroma active compounds. *Meat Science*, 93, 776–785, <http://dx.doi.org/10.1016/j.meatsci.2012.11.040>
- Desmond, E. (2006). Reducing salt: A challenge for the meat industry. *Meat Science*, 74, 188–196, <http://dx.doi.org/10.1016/j.meatsci.2006.04.014>
- Doyon, M. & Labrecque, J. (2008). Functional foods : A conceptual definition. *British Food Journal*, 110, 1133–1149, <http://dx.doi.org/10.1108/00070700810918036>
- Galanakis, C. M., Rizou, M., Aldawoud, T. M. S., Ucak, I. & Rowan, N. J. (2021). Innovations and technology disruptions in the food sector within the COVID-19 pandemic and post-lockdown era. *Trends in Food Science & Technology*, 110, 193–200, <https://doi.org/10.1016/j.tifs.2021.02.002>
- Grasso, S. (2020). Hybrid meat. *Food Science and Technology*, 34, 48–51, [https://doi.org/10.1002/fsat.3403\\_12.x](https://doi.org/10.1002/fsat.3403_12.x)
- Grasso, S. & Jaworska, S. (2020). Part meat and part plant: Are hybrid meat products fad or future? *Foods*, 9, 1888, <https://doi.org/10.3390%2Ffoods9121888>
- Heck, R. T., Fagundes, M. B., Cichoski, A. J., de Menezes, C. R., Barin, J. S., Lorenzo, J. M. . . . & Campagnol, P. C. B. (2019). Volatile compounds and sensory profile of burgers with 50% fat replacement by microparticles of chia oil enriched with rosemary. *Meat Science*, 148, 164–170, <https://doi.org/10.1016/j.meatsci.2018.10.017>
- Heck, R. T., Vendruscolo, R. G., de Araújo Etchepare, M., Cichoski, A. J., de Menezes, C. R., Barin, J. S. . . . & Campagnol, P. C. B. (2017). Is it possible to produce a low-fat burger with a healthy n-6/n-3 PUFA ratio without affecting the technological and sensory properties? *Meat Science*, 130, 16–25, <https://doi.org/10.1016/j.meatsci.2017.03.010>
- Inguglia, E. S., Zhang, Z., Tiwari, B. K., Kerry, J. P. & Burgess, C. M. (2017). Salt reduction strategies in processed meat products — A review. *Trends in Food Science & Technology*, 59, 70–78, <https://doi.org/10.1016/j.tifs.2016.10.016>
- Iwatani, S. & Yamamoto, N. (2019). Functional food products in Japan: A review. *Food Science and Human Wellness*, 8, 96–101, <https://doi.org/10.1016/j.fshw.2019.03.011>
- Jiménez-Colmenero, F., Carballo, J. & Cofrades, S. (2001). Healthier meat and meat products: their role as functional foods. *Meat Science*, 59, 5–13, [http://dx.doi.org/10.1016/S0309-1740\(01\)00053-5](http://dx.doi.org/10.1016/S0309-1740(01)00053-5)
- Josquin, N. M., Linssen, J. P. H. & Houben, J. H. (2012). Quality characteristics of Dutch-style fermented sausages manufactured with partial replacement of pork back-fat with pure, pre-emulsified or encapsulated fish oil. *Meat Science*, 90, 81–86, <http://dx.doi.org/10.1016/j.meatsci.2011.06.001>
- Kurčubić, V., Okanović, D., Vasilev, D., Ivić, M., Čolović, D., Jokanović, M. & Džinić, N. (2020). Effects of replacing pork back fat with cellulose fiber in Pariser sausages. *Fleischwirtschaft*, 100, 82–88.
- Kurčubić, V., Stajić, S., Miletić, N. & Stanišić, N. (2022). Healthier meat products are fashionable — consumers love fashion. *Applied Sciences*, 12, 10129, <https://doi.org/10.3390/app121910129>
- Kurčubić, V. S., Mašković, P. Z., Vujić, J. M., Vranić, D. V., Vesković-Moračanin, S. M., Okanović, Đ. G. & Lilić, S. V. (2014). Antioxidant and antimicrobial activity of *Kitabelia vitifolia* extract as alternative to the added nitrite in fermented dry sausage. *Meat Science*, 97, 459–467, <http://dx.doi.org/10.1016/j.meatsci.2014.03.012>
- Lawrie, R. A. & Ledward, D. (2006). *Lawrie's Meat Science, Seventh Edition*: CRC Press.
- Liaros, N. G., Katsanidis, E. & Bloukas, J. G. (2009). Effect of the ripening time under vacuum and packaging film permeability on processing and quality characteristics of low-fat fermented sausages. *Meat Science*, 83, 589–598, <http://dx.doi.org/10.1016/j.meatsci.2009.07.006>

- Olmedilla-Alonso, B., Jiménez-Colmenero, F. & Sánchez-Muniz, F. J. (2013).** Development and assessment of healthy properties of meat and meat products designed as functional foods. *Meat Science*, 95, 919–930, <http://dx.doi.org/10.1016/j.meatsci.2013.03.030>
- Pereira, P. M. & Vicente, A. F. (2013).** Meat nutritional composition and nutritive role in the human diet. *Meat Science*, 93, 586–592, <http://dx.doi.org/10.1016/j.meatsci.2012.09.018>
- Petit, G., Jury, V., de Lamballerie, M., Durantou, F., Potier, L. & Martín, J.-L. (2019).** Salt intake from processed meat products: Benefits, risks and evolving practices. *Comprehensive Reviews in Food Science and Food Safety*, 18, 1453–1473, <https://doi.org/10.1111/1541-4337.12478>
- Pintado, T. & Cofrades, S. (2020).** Quality characteristics of healthy dry fermented sausages formulated with a mixture of olive and chia oil structured in oleogel or emulsion gel as animal fat replacer. *Foods*, 9, 830.
- Pinton, M. B., Correa, L. P., Facchi, M. M. X., Heck, R. T., Leães, Y. S. V., Cichoski, A. J. . . . & Campagnol, P. C. B. (2019).** Ultrasound: A new approach to reduce phosphate content of meat emulsions. *Meat Science*, 152, 88–95, <https://doi.org/10.1016/j.meatsci.2019.02.010>
- Powell, M. J., Sebranek, J. G., Prusa, K. J. & Tarté, R. (2019).** Evaluation of citrus fiber as a natural replacer of sodium phosphate in alternatively-cured all-pork Bologna sausage. *Meat Science*, 157, 107883, <https://doi.org/10.1016/j.meatsci.2019.107883>
- Sebranek, J. G. (2009).** Basic Curing Ingredients. In R. Tarté (Ed.), *Ingredients in Meat Products: Properties, Functionality and Applications* (pp. 1–23). New York, NY: Springer, New York.
- Sindelar, J. J. & Milkowski, A. L. (2012).** Human safety controversies surrounding nitrate and nitrite in the diet. *Nitric Oxide*, 26, 259–266, <http://dx.doi.org/10.1016/j.niox.2012.03.011>
- Singh, P., Kumar, R., Sabapathy, S. & Bawa, A. (2008).** Functional and edible uses of soy protein products. *Comprehensive Reviews in Food Science and Food Safety*, 7, 14–28, <https://doi.org/10.1111/j.1541-4337.2007.00025.x>
- Siró, I., Kápolna, E., Kápolna, B. & Lugasi, A. (2008).** Functional food. Product development, marketing and consumer acceptance — A review. *Appetite*, 51, 456–467, <https://doi.org/10.1016/j.appet.2008.05.060>
- Stajić, S., Kalušević, A., Tomasevic, I., Rabrenović, B., Božić, A., Radović, P. . . . & Živković, D. (2020).** Technological properties of model system beef emulsions with encapsulated pumpkin seed oil and shell powder. *Polish Journal of Food and Nutrition Sciences*, 70(2), 159–168, <https://doi.org/10.31883/pjfn/118008>
- Stajić, S., Lilić, S., Danijela, V., Vladimir Tomović & Živković, D. (2018).** *Sastav lipida fermentisane kobasica sa dodatkom lanenog ulja*. Paper presented at the XXI-II Savetovanje o biotehnologiji, Čačak, Srbija.
- Stajić, S., Stanišić, N., Tomasevic, I., Djekic, I., Ivanović, N. & Živković, D. (2018).** Use of linseed oil in improving the quality of chicken frankfurters. *Journal of Food Processing and Preservation*, 42, e13529, <http://doi.org/10.1111/jfpp.13529>
- Stajić, S., Tomasevic, I., Stanišić, N., Tomović, V., Lilić, S., Vranić, D. . . . & Živković, D. (2020).** Quality of dry-fermented sausages with backfat replacement — Fermented sausages with high content of flaxseed oil and pre-treated with soy protein isolate and alginate. *Fleischwirtschaft*, 100(7), 74–81.
- Stajić, S., Tomasevic, I., Tomovic, V. & Stanišić, N. (2022).** Dietary fibre as phosphate replacement in all-beef model system emulsions with reduced content of sodium chloride. *Journal of Food and Nutrition Research*, 61, 277–285.
- Stajić, S. & Vasilev, D. (2022).** Encapsulation of meat products ingredients and influence on product quality. In S. Lević, V. Nedović & B. Bugarski (Eds.), *Encapsulation in Food Processing and Fermentation*. Boca Raton: CRC Press.
- Talens, C., Ibagüen, M., Murgui, X., García-Muñoz, S. & Peral, I. (2022).** Texture-modified meat for senior consumers varying meat type and mincing speed: Effect of gender, age and nutritional information on sensory perception and preferences. *Future Foods*, 6, 100180, <https://doi.org/10.1016/j.fufo.2022.100180>
- Tungland, B. C. & Meyer, D. (2002).** Nondigestible oligo- and polysaccharides (dietary fiber): Their physiology and role in human health and food. *Comprehensive Reviews in Food Science and Food Safety*, 1, 90–109, <https://doi.org/10.1111/j.1541-4337.2002.tb00009.x>
- Vasquez Mejia, S. M., Shaheen, A., Zhou, Z., McNeill, D. & Bohrer, B. M. (2019).** The effect of specialty salts on cooking loss, texture properties, and instrumental color of beef emulsion modeling systems. *Meat Science*, 156, 85–92, <https://doi.org/10.1016/j.meatsci.2019.05.015>
- Williams, P. (2007).** Nutritional composition of red meat. *Nutrition & Dietetics*, 64, S113–S119, <https://doi.org/10.1111/j.1747-0080.2007.00197.x>
- Wood, J. D., Enser, M., Fisher, A. V., Nute, G. R., Sheard, P. R., Richardson, R. I. . . . & Whittington, F. M. (2008).** Fat deposition, fatty acid composition and meat quality: A review. *Meat Science*, 78(4), 343–358, <http://dx.doi.org/10.1016/j.meatsci.2007.07.019>
- Young, J. F., Therkildsen, M., Ekstrand, B., Che, B. N., Larsen, M. K., Oksbjerg, N. & Stagsted, J. (2013).** Novel aspects of health promoting compounds in meat. *Meat Science*, 95, 904–911, <http://dx.doi.org/10.1016/j.meatsci.2013.04.036>
- Zhang, W., Xiao, S., Samaraweera, H., Lee, E. J. & Ahn, D. U. (2010).** Improving functional value of meat products. *Meat Science*, 86, 15–31, <http://dx.doi.org/10.1016/j.meatsci.2010.04.018>