



Honeybee pollen as a bioindicator of contamination: an overview

Jelena Ćirić^{a*}, Nils Haneklaus^{b,c}, Tatjana Baltić^a, Sara Simunović^a, Nenad Parunović^a,
Dejana Trbović^a and Boris Mrdović^a

^a Institute of Meat Hygiene and Technology, Kačanskog 13, 11040 Belgrade, Serbia

^b Universität für Weiterbildung Krems, 3500, Krems an der Donau, Austria

^c Technische Universität Bergakademie Freiberg, 09599, Freiberg, Germany

ARTICLE INFO

Keywords:

Apiculture
Bee pollen
Pesticide
Heavy metal
Residue

ABSTRACT

Honeybees and honeybee products (honey, bee wax, bee pollen and bee bread) are potential bioindicators of contaminants (pesticides, mycotoxins, pyrrolizidine alkaloids, toxic elements, radionuclides etc.) in the environment. In this study, recent results on the food safety risks of bee pollen and data about the concentration of toxic substances detected in bee pollen are summarized. Based on different studies, a risk assessment was conducted for the most common pesticide active substances (chlorpyrifos, fluralanone, carbendazim, thiacloprid), heavy metals (arsenic, cadmium, mercury, lead) and common mycotoxins (aflatoxin-B1, ochratoxin-A, fumonisins, zearalenone, deoxynivalenol, T-2 toxin).

1. Introduction

Bee pollen contains essential nutrients. According to Campos *et al.*, (2008), carbohydrates (13–55%), proteins (10–40%), lipids (1–13%) and fibre (0.3–20%) all contribute to the composition of bee pollen. In addition, bee pollens are rich in biologically active micronutrients like minerals, polyphenols and vitamins. Based on a report by Habryka *et al.*, (2016), the product is used in apitherapy mainly for its antioxidant, anti-inflammatory, anti-biotic and antiallergic effects. Bee pollen improves blood supply to the nerve tissue, thereby increasing mental performance and eliminating the state of fatigue. Research works have also shown a positive effect of bee pollen on some diseases of the liver, heart and prostate. The main consumers of bee pollen are the followers of health- and environmental-

ly-conscious lifestyles, as well as the elderly, who use it due to its antioxidant and other therapeutic effects (Végh *et al.*, 2021).

Besides all of these positive effects and high biological and nutritive value, bee pollen can contain hazardous trace elements, pesticide residues (Ambrus *et al.*, 2020; Bostan *et al.*, 2019; Toselli and Sgolastra, 2020), toxic metals and metalloids (Spirić *et al.*, 2019; Ćirić *et al.*, 2021; Murashova *et al.*, 2020; Roman, 2009), moulds and mycotoxins (Alarcón *et al.*, 2019), pyrrolizidine alkaloids (Botías *et al.*, 2015), allergens (Pitsios *et al.*, 2006; Nonotte-Varly, 2016) and GM (genetically modified) foods (Malone, 2002). To ensure the safety and quality of bee pollen, some countries issued national legislations, decisions and guidelines which correlate with European and International standards. In Serbia for instance, honeybee

*Corresponding author: Jelena Ćirić, jelena.ciric@inmes.rs

Paper received Jun 6th 2023. Paper accepted Jun 18th 2023.

Published by Institute of Meat Hygiene and Technology — Belgrade, Serbia

This is an open access article under CC BY licence (<http://creativecommons.org/licenses/by/4.0>)

Table 1. Pesticide residue contents of bee pollen from different studies

Pesticide Concentration Mean value	Active ingredients	Country of origin	Reference
30 µg/kg 61 µg/kg 16 µg/kg	Tebuconazole Thiacloprid Chlorpyrifos	Poland	<i>Roszko et al. (2016)</i>
470 µg/kg 79 µg/kg	Thiacloprid Prothioconazol-desthio	Germany	<i>Böhme et al. (2018)</i>
133 µg/kg 40 µg/kg	Thiacloprid Permethrin-cis	Luxembourg	<i>Beyer et al. (2018)</i>
24 µg/kg 7 µg/kg	Carbendazim Amitraz II	France	<i>Lambert et al. (2013)</i>
915 µg/kg 128 µg/kg 83 µg/kg	Fluvalinate Chlorpyrifos Carbaryl	Taiwan	<i>Nai et al. (2017)</i>
6 µg/kg 28 µg/kg 227 µg/kg 50 µg/kg 3 µg/kg	Coumaphos Carbaryl Phosmet Carbendazim Atrazine	USA	<i>Stoner and Eitzer (2013)</i>
1 µg/kg 3 µg/kg <1 µg/kg	Azoxystrobin Carbendazim Carbaryl	Uruguay	<i>Niell et al. (2015)</i>

products must meet legal criteria (*Republic of Serbia*, 2015). Maximum Residue Level (MRL) values for honey vary between 0.01 and 1 mg/kg, but for other honeybee products, no MRLs are applicable until individual products have been identified and listed (*EU Pesticides Database*, 2021). The website of the International Honey Commission (IHC) is often quoted in scientific research in which maximum limits have been proposed for Pb (500 µg/kg), Cd (30 µg/kg) and Hg (10 µg/kg) in honey. Applying the data of Table 1, risk assessments for

pesticide residues in bee pollen were performed in different studies. The results indicate that a major pesticide is thiacloprid.

Table 2 summarizes the literature data on the concentration of toxicologically important elements (As, Cd, Hg, Pb) in bee pollen. The mean values for Cd concentration of bee pollen samples exceeded 30 µg/kg in most studies, except in Brazil (*de Oliveira et al.*, 2017). In some bee pollens from Europe, the Pb concentrations exceeded the 200 µg/kg limit (*Lambert et al. 2012; Adaškevičiūtė et al.*, 2019).

Table 2. Toxic metal contamination of bee pollen from different studies

Mean concentration of toxic elements (µg/kg)				Country of origin	Reference
Lead (Pb)	Arsenic (As)	Mercury (Hg)	Cadmium (Cd)		
112	/	/	30	Italy	<i>Conti and Botré (2001)</i>
237	/	/	/	France	<i>Lambert et al. (2012)</i>
247	/	/	88	Europe	<i>Adaškevičiūtė et al. (2019)</i>
20	/	/	20	Chile	<i>Mejías et al. (2018)</i>
148	169	/	2	Brazil	<i>de Oliveira et al. (2017)</i>

Table 3. Mould contamination of bee pollen from different studies

Isolated mould	Country of origin	Reference
Alternaria Aspergillus Fusarium Mucor Penicillium Rhizopus	Serbia	<i>Kostić et al.</i> (2017)
Alternaria Aspergillus Cladosporium Penicillium Rhizopus	Ukraine, Slovakia	<i>Shevtsova et al.</i> (2014)
Alternaria Aspergillus Cladosporium Fusarium Mucor Paecilomyces Penicillium Rhizopus	Slovakia	<i>Kačániová et al.</i> (2011)

Over the past decades, several studies have been conducted in Europe on the mycotoxin content of bee pollen samples. The results of the reviewed studies are presented in Table 3. *Kostić et al.* (2017)

reported that the average concentrations of aflatoxin-B1 exceeded the legal limit by more than four times in multifloral pollen samples from Serbia. The same authors isolated different moulds from bee pollen in Serbia: *Alternaria*, *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium* and *Rhizopus*. Also, similar moulds were isolated in a study from Slovakia and Ukraine (*Shevtsova et al.*, 2014).

2. Conclusions

Bee pollen is a very popular bee product that is presently not defined in most national regulations. In Europe, many studies have been conducted on this bee product, but little data is available from other continents. According to different studies, bee pollen is characterized by heterogenous food safety risks and could also be used as a potential environmental bioindicator. The common contaminants of bee pollen are pesticides, heavy metals, metalloids and mycotoxins. In this short overview, recent findings on the above-mentioned substances and data on concentrations determined in bee pollen were summarized, from different countries. A number of scientific works on the topic of bee pollen is associated with biomonitoring systems, and bee pollen can indeed be used as an environmental bioindicator, similar to the way other bee products (bee wax, honey and bee bread) are already in use.

Disclosure statement: No potential conflict of interest was reported by the authors.

Funding: This study was supported by the Ministry of Science, Technological Development and Innovation, Republic of Serbia, Grant No. 451-03-47/2023-01/200050 from 03.02.2023 and the German Federal Ministry of Education and Research (BMBF) (Bridge2ERA2021: 100579052).

References

- Adaškevičiūtė, V., Kaškonienė, V., Kaškonas, P., Barčauskaitė, K. & Maruška, A. (2019). Comparison of physicochemical properties of bee pollen with other bee products. *Biomolecules*, 9(12), 819.
- Alarcón, D. C. M., Cáceres, E. X. U. & Becerra, A. M. A. (2019). Mycotoxins in foods that cause damage to humans. *Journal of Applied Biotechnology & Bioengineering*, 6, 259–263.
- Ambrus, Á., Szenczi-Cseh, J., Griff, T., Kerekes, K., Miklós, G., Vásárhelyi A. & Szigeti, T. J. (2020). Food safety assessment of the mycotoxin and pesticide residue contamination of our foods, Part 2. Mycotoxins. *Journal of Food Investigation*, 66, 2938–2949.
- Beyer, M., Lenouvel, A., Guignard, C., Eickermann, M., Clermont, A., Kraus, F. & Hoffmann, L. (2018). Pesticide residue profiles in bee bread and pollen samples and the survival of honeybee colonies—a case study from Luxembourg. *Environmental Science and Pollution Research*, 25, 32163–32177.
- Böhme, F., Bischoff, G., Zebitz, C. P., Rosenkranz, P. & Wallner, K. (2018). Pesticide residue survey of pollen loads collected by honeybees (*Apis mellifera*) in daily intervals at three agricultural sites in South Germany. *PLoS One*, 13(7), e0199995.

- Bostan, I., Onofrei, M., Gavriluță, A. F., Toderășcu, C. & Lazăr, C. M. (2019). An integrated approach to current trends in organic food in the EU. *Foods*, 8(5), 144.
- Bofías, C., David, A., Horwood, J., Abdul-Sada, A., Nicholls, E., Hill, E. & Goulson, D. (2015). Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. *Environmental Science & Technology*, 49(21), 12731–12740.
- Campos, M. G., Bogdanov, S., de Almeida-Muradian, L. B., Szczesna, T., Mancebo, Y., Frigerio, C. & Ferreira, F. (2008). Pollen composition and standardisation of analytical methods. *Journal of Apicultural Research*, 47(2), 154–161.
- Conti, M. E. & Botrè, F. (2001). Honeybees and their products as potential bioindicators of heavy metals contamination. *Environmental Monitoring and Assessment*, 69, 267–282.
- Ćirić, J., Spirić, D., Baltić, T., Lazić, I. B., Trbović, D., Parunović, N., Petronijević, R. & Đorđević V., 2020. Honey bees and their products as indicators of environmental element deposition. *Biological Trace Element Research*, 199, 2312–2319, doi: 10.1007/s12011-020-02321-6. Epub 2020 Aug 5. PMID: 32757175
- de Oliveira, F. A., de Abreu, A. T., de Oliveira Nascimento, N., Froes-Silva, R. E. S., Antonini, Y., Nalini Jr, H. A. & de Lena, J. C. (2017). Evaluation of matrix effect on the determination of rare earth elements and As, Bi, Cd, Pb, Se and In in honey and pollen of native Brazilian bees (*Tetragonisca angustula*-Jataí) by Q-ICP-MS. *Talanta*, 162, 488–494.
- EU Pesticides Database, (2021), <https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activesubstance.selection&language=EN/> (2021), Accessed 19th Mar 2021.
- Habryka, C., Kruczek, M. & Drygaś, B. (2016). Bee products used in apitherapy. *World Scientific News*, 48, 254–258.
- Kačaniová, M., Juráček, M., Chlebo, R., Kňazovická, V., Kadasi-Horáková, M., Kunová, S., ... & Šimko, M. (2011). Mycobiota and mycotoxins in bee pollen collected from different areas of Slovakia. *Journal of Environmental Science and Health, Part B*, 46(7), 623–629.
- Kostić, A. Ž., Petrović, T. S., Krnjaja, V. S., Nedić, N. M., Tešić, Ž. L., Milojković-Opsenica, D. M., ... & Pešić, M. B. (2017). Mold/aflatoxin contamination of honey bee collected pollen from different Serbian regions. *Journal of Apicultural Research*, 56(1), 13–20.
- Lambert, O., Piroux, M., Puyo, S., Thorin, C., L'Hostis, M., Wiest, L., ... & Pouliquen, H. (2013). Widespread occurrence of chemical residues in beehive matrices from apiaries located in different landscapes of Western France. *PLoS one*, 8(6), e67007.
- Malone, L. A. (2002). Literature review on genetically modified plants and bee products. HortResearch.
- Mejías, E., Gómez, C. J., Gareil, P., Delaunay, N. & Montenegro, G. (2018). Characterization of phenolic profile alterations in metal-polluted bee pollen via capillary electrophoresis. *Ciencia e Investigación Agraria: Revista Latinoamericana de Ciencias de la Agricultura*, 45(1), 51–63.
- Murashova, E. A., Tunikov, G. M., Nefedova, S. A., Karelina, O. A., Byshova, N. G. & Serebryakova, O. V. (2020). Major factors determining accumulation of toxic elements by bees and honey products. *International Transaction Journal of Engineering, Management and Applied Sciences and Technologies*, 11(3), 11A03N.
- Nai, Y. S., Chen, T. Y., Chen, Y. C., Chen, C. T., Chen, B. Y. & Chen, Y. W. (2017). Revealing pesticide residues under high pesticide stress in Taiwan's agricultural environment probed by fresh honey bee (Hymenoptera: Apidae) Pollen. *Journal of Economic Entomology*, 110(5), 1947–1958.
- Niell, S., Jesus, F., Pérez, C., Mendoza, Y., Diaz, R., Franco, J., ... & Heinzen, H. (2015). QuEChERS adaptability for the analysis of pesticide residues in beehive products seeking the development of an agroecosystem sustainability monitor. *Journal of Agricultural and Food Chemistry*, 63(18), 4484–4492.
- Nonotte-Varly, C. (2016). Why not evaluate the allergenic potential of bee pollen with a skin testing method? *Alergologia Polska [Polish Journal of Allergology]*, 3(3), 114–122.
- Pitsios, C., Chliva, C., Mikos, N., Kompoti, E., Nowak-Węgrzyn, A. & Kontou-Fili, K. (2006). Bee pollen sensitivity in airborne pollen allergic individuals. *Annals of Allergy, Asthma & Immunology*, 97(5), 703–706.
- Republic of Serbia, (2015). Rulebook on Quality of Honey and Other Bee Products. *Official Gazette of the Republic of Serbia*, 101.
- Roman, A. (2009). Concentration of chosen trace elements of toxic properties in bee pollen loads. *Polish Journal of Environmental Studies*, 18(2).
- Roszkó, M. Ł., Kamińska, M., Szymczyk, K. & Jędrzejczak, R. (2016). Levels of selected persistent organic pollutants (PCB, PBDE) and pesticides in honey bee pollen sampled in Poland. *PLoS One*, 11(12), e0167487.
- Shevtsova, T., Kačaniová, M., Garkava, K., Brindza, J. & Petrova, J. (2014). Contamination of *Betula verrucosa* Ehrh. pollen by microorganisms, mycotoxins and heavy metals, https://dspace.nau.edu.ua/bitstream/NAU/10121/1/jmbfs_xxs1_shevtsova.pdf
- Spirić, D., Ćirić, J., Đorđević, V., Nikolić, D., Janković, S., Nikolić, A... & Teodorović, V. (2019). Toxic and essential element concentrations in different honey types. *International Journal of Environmental Analytical Chemistry*, 99(5), 474–485.
- Stoner, K. A. & Eitzer, B. D. (2013). Using a hazard quotient to evaluate pesticide residues detected in pollen trapped from honey bees (*Apis mellifera*) in Connecticut. *PLoS One*, 8(10), e77550.
- Toselli, G. & Sgolastra, F. (2020). Seek and you shall find: An assessment of the influence of the analytical methodologies on pesticide occurrences in honey bee-collected pollen with a systematic review. *Chemosphere*, 258, 127358.
- Végh, R., Csóka, M., Sörös, C. & Sipos, L. (2021). Food safety hazards of bee pollen—A review. *Trends in Food Science & Technology*, 114, 490–509.